

UNCUT-NEWS



[Demokratizemedia CC0](#)

Kennedy reveals the shocking truth about vaccines in 3 minutes

- uncut-news.ch
- [November 15, 2024](#)

Robert F. Kennedy Jr. called Anthony Fauci a liar when he claimed that none of the 72 childhood vaccines in the United States had ever been tested for safety. After a year-long legal battle, Fauci's lawyers admitted that Kennedy was right.



Wide Awake Media 
@wideawake_media · [Follow](#)



Having been called a liar by Anthony Fauci for saying that "not one of the 72 vaccines mandated for children has ever been safety tested", RFK Jr. sued Fauci.

After a year of stonewalling, Fauci's lawyers admitted that RFK Jr. had been right all along.

"There's no downstream... [Mehr anzeigen](#)



11:41 a.m. · November 12 2024



Kennedy says vaccine manufacturers can't be held accountable, their products aren't tested, and they don't invest in marketing because the government requires 78 million school children to be vaccinated every year. In a speech at Hillsdale College, Kennedy said this was a perfect business model for manufacturers. As soon as a vaccine is included in the program, the company will generate annual revenue of one billion dollars.

There are currently 72 approved childhood vaccines in the U.S. and 16 others. The Vaccine Injury Act of 1986 exempted vaccine manufacturers from liability. According to Kennedy, three years later, in 1989, cases of chronic diseases among American children skyrocketed. He cited conditions such as ADHD, sleep disorders, speech problems, autism, Tourette syndrome and narcolepsy.

Increasing numbers and new plans

"The rate of autism went from one case in 10,000 children in my generation to one in 34 children today."

Donald Trump plans to give Kennedy a major role in public health because, according to Trump, he knows "more than anyone" about health care.

Not a single routine childhood vaccine has been approved on the basis of a long-term placebo-controlled trial. Not a single one.



Aaron Siri
@AaronSiriSG · Follow



Not a single routine childhood vaccine was licensed based on a long-term placebo-controlled trial. Not one. Thank you @ICANdecide for funding the careful creation of this chart. icandecide.org/no-placebo



None of the vaccine doses the CDC recommends for routine injection into children were licensed by the FDA based on a long-term placebo-controlled trial

Type	Doses	Age Injected	Brand	Company	Control	Placebo?	Safety Review After Injection	Long	Source	Note
HepB	3	Birth 1M 6M	Recombivax HB	M	None	NO	5 days	NO	Package insert at § 6.1	Note that to license a vaccine for children, the FDA relies upon the clinical trial conducted with children, not adults, because as the FDA explains, "It's important that the public recognize that, because young children are still growing and developing, it's critical that thorough and robust clinical trials of adequate size are completed to evaluate the safety and the immune response to a vaccine in this population. Children are not small adults."
			Engerix B	G	None	NO	4 days	NO	Package insert at § 6.1	
DTaP	15	2M 4M/6M 15M 4Y	Infanrix	G	DTP	NO	30 days	NO	Package insert at § 6.1	DTP was also not licensed based on a placebo controlled trial and it increases mortality .
			Daptacel	S	DT or DTP	NO	Up to 2 months + 1 trial 6 months	NO	Package insert at § 6.1	The 6-month Daptacel trial had no control, 1,454 children and "[w]ithin 30 days following any dose of DAPTACEL, 3.9% subjects reported at least one serious adverse event ."
PCV	4	2M 4M 6M 12M	Prenar 13, PCV-13	P	Prenar 7	NO	6 months	NO	Package insert at § 6.1	Prenar 7 trial's control was an "[i]nvestigational meningococcal group C conjugate vaccine." In Prenar 13 trial, "[s]erious adverse events reported following vaccination in infants and toddlers occurred in 8.2% among Prenar 13 recipients and 7.2% among Prenar 7 recipients." In Vaxneuvance trial, "serious adverse events... were reported by 9.6% of VAXNEUVANCE recipients and by 8.9% of Prenar 13 recipients" but deemed "safe" because "no notable patterns or numerical imbalances between vaccination groups." Prenar 20 had similar result split into "serious adverse events" and "newly diagnosed chronic medical conditions."
			Vaxneuvance PCV-15	M	Prenar 13	NO	6 months	NO	Package insert at § 6.1	
			Prenar 20, PCV-20	P	Prenar 13	NO	6 months	NO	Package insert at § 6.1; Clinical Review	
IPV	4	2M/4M 6M/4Y	IPOL	S	None	NO	3 days	NO	Package insert at 14-17	IPOL is very different than the polio vaccine created by Jonas Salk in the 1950s (used until 1960s). Hence, trials of Salk's vaccine from the 1950s were not relied upon to license IPOL.
Hib	3 or 4	2M 4M 6M 12M	ActHib	S	HepB	NO	30 days	NO	Package insert at § 6.1; Letter of Approval at 8	Within 30 days of injection in the ActHib trial, 3.4% experienced a serious adverse event but "[n]one was assessed by the investigators [Sonaf] as related to the study of vaccines."
			Hiberix	G	HibTITER or other vaccine	NO	31 days	NO	Package insert at § 6.1; Clinical Review at 20-21	
			Liquid PedvaxHib	M	Lyophilized PedvaxHib	NO	3 days	NO	Package insert at 6-8	Lyophilized PedvaxHib vaccine, used as the control for Liquid PedvaxHib, was tested in a trial in which controls were given placebo, OPV, and DTP but there is no indication Lyophilized PedvaxHib was ever licensed.
RV	2 or 3	2M 4M 6M	Rotarix	G	None	NO	31 days + 1 year for intubation	NO	Package insert at § 6.1; Clinical Review at 23-24	"[T]here were 68 (0.19%) deaths following ROTARIX and 50 (0.15%) deaths following placebo... The most common cause... was pneumonia... observed in 29 (0.05%) recipients of ROTARIX and 10 (0.03%) placebo recipients." Its clinical review admits "[t]he placebo consisted of all components of Rotarix, but without any RV particles." The package insert for RotaTeq similarly admits its "placebo" contains multiple ingredients as seen to the left.
			RotaTeq	M	None	NO	42 days + 1 year for intubation	NO	Package insert at § 6.1; Clinical Review at 445 etc.	
Covid19	3	6M 7M 10M	Comirnaty	P	Placebo	YES	6 months	NO	Package insert at § 6.1	Comirnaty licensed for only 12+ (Spikevax, Moderna, only 18+). Placebo controls unblinded and most vaccinated during the trial. All data 16+ is combined but 12-15 data is separate, had 1,131 vaccinated children, and one participant shows how this trial was conducted.
Flu	19	6M 7M Yearly	Various	Various	Flu shots change annually without any clinical trial	NO	Flu shots change annually without any clinical trial	NO	CDC 22-23 Flu Shots; FDA Flu Shots	The trials of the original flu shot formulations for children also did not have a placebo control (see pp. 13-14) even though some adult trials did. The one inhaled influenza vaccine had a placebo but, again, it changes every year and is not safety tested in any trial.
MMR	6	12M 4Y	M-M-R-II	M	None	NO	42 days	NO	Clinical reports	M-M-R-II trials totaled only 834 children and a third developed gastrointestinal issues and a third respiratory issues. In Priorix trial, both vaccine groups had high rate of serious adverse events, emergency room visits, and new chronic diseases (e.g., autoimmune disorders, asthma, type I diabetes, celiac, and allergies). See Table 6 of the Supplementary Materials.
			Priorix	G	M-M-R-II	NO	6 months	NO	Package insert at § 6.1; Sup materials at 12	
VAR	2	12M 4Y	Varivax	M	45 mg of neomycin per milliliter	NO	70 days	NO	Package insert at § 6.1; Merck study at 2; Clinical reports	One controlled trial with 956 children, half Varivax and half neomycin, and one trial with 32 vaccinated and another 29 vaccinated 8 weeks later, during which the first group had double the ear infections and 50% more respiratory infections.
HepA	2	12M 18M	Havrix	G	Engerix-B	NO	6 months	NO	Package insert at § 6.1	Trials for both occurred at the same time when there was no licensed Hep A vaccine and hence no excuse for not using a placebo control. It is also startling Engerix-B, see above, was the control for Havrix, and an injection of cyto-and-neuro toxic substances, AAHS and thimerosal, were used as a control for Varivax instead of a saline injection.
			Vagta	M	AAHS and Thimerosal	NO	42 days	NO	Package insert at § 6.1; Merck study at 454	
Tdap	3	11Y	Adacel	S	Td, for adults	NO	6 months	NO	Package insert at § 6.1	Due to reactions, Tdap (Adacel) given at 11Y has 12.5 times less diphtheria toxin (25U v 2U) and 10 times less pertussis toxin (25mcg v 2.5mcg) than DTaP (Infanrix) given to babies.
			Boostrix	G	Denarix or Adacel	NO	6 months	NO	Package insert at § 6.1	
HPV	2 or 3	9Y 9Y 9Y	Gardasil 9	M	Gardasil 4 (see note)	NO	1 month in five trials, 6 months in one trial, and 4 years in one trial	NO	Clinical review at 17-19	Gardasil 9 trial gave 306 people placebo after full series of Gardasil 4. In Gardasil 4's trial , controls received aluminum adjuvant, AAHS, except 320 people labeled "Saline-Placebo" that actually received all vaccine ingredients except antigens and AAHS. Across trials, 2-3% receiving vaccine or aluminum adjuvant (used to induce autoimmunity) had a suspected autoimmune disorder.
			Menactra	S	Menomune	NO	6 months	NO	Package insert at § 6.1	Incredibly, the safety section of the package insert for Menomune lists the trial in which it was used as a control for the trial of Menactra. This provides another good example of the safety pyramid scheme in which Menomune is licensed without a placebo-controlled trial and then used as the control to license Menactra; Menactra is then used as the control to license
Men4	2	11Y	Menveo	G	Menactra or	NO	6 months	NO	Package insert at § 6.1	

7:50 p.m. · July 23, 2024

