

Shocking “Sudden and Unexpected” study – How deadly are the COVID vaccines actually?

- 27. August 2024



Picture: freepik / user7350813

Once again, a study on the fatal side effects of experimental COVID vaccines causes a stir. According to Jordanian data, three percent of vaccinators died "suddenly and unexpectedly" shortly after the vaccine was administered. Newcomers to the world, this would correspond to around 150 million vaccination deaths...

A recently published study by Jordanian scientists has attracted attention by investigating possible relationships between COVID-19 vaccinations and unexpected health consequences. The research, conducted by Nadia Al-Rousan and Hazem Al-Najjar from the German-Jordanian University of SEEIT and the University of Petra, analysed data on side effects after COVID-19 vaccinations.

“Het vaccinatiegerelateerde sterftecijfer bedraagt ongeveer 3% van degenen die het vaccin hebben gekregen”

Evaluation of the effects of MERCK, MODERNA, PFIZER/BioNTech, and JANSSEN COVID-19 vaccines on vaccinated people: A metadata analysis <https://t.co/LJViQH4jDx>
[pic.twitter.com/ak9kukIFMB](https://t.co/LJViQH4jDx)

Lilian (Lily4Lily2) [August 26, 2024](#)

The study entitled “[MERCK, MODERNA, PFIZER/BioNTech, and JANSSEN COVID-19 vaccines on vaccinated people: A metadata analysis](#)” was published in the journal

“Informatics in Medicine Unlocked”. It is based on the evaluation of approximately 65,000 cases documented by the World Health Organisation (WHO).

Evaluation of the effects of MERCK, MODERNA, PFIZER/BioNTech, and JANSSEN COVID-19 vaccines on vaccinated people: A metadata analysis <https://t.co/LplX6swSdy>
[pic.twitter.com/PtTgjKoLD1](https://t.co/LplX6swSdy)

Sabine Stebel (Sabisteb) [August 26, 2024](#)

The central statement of the researchers is:

“The overall mortality rate in the vaccinated population is remarkable. 40 different mild to severe symptoms were reported in the vaccinated individuals. The study highlights the 10 most common symptoms that occurred after vaccination. Women under 60 account for the majority of the data set.”

One of the most striking conclusions is:

“The vaccine-related mortality rate is about 3 percent of the vaccinated, with most cases occurring in people under 60 who were not hospitalized and had received their first dose of vaccination.”

The study identified headaches, shivering and fatigue as the three most commonly reported symptoms after vaccination. Particularly worrying is the finding that 97.4 percent of registered deaths affected persons under the age of 60. The authors emphasize the importance of the first days after the vaccination: “The first three days after vaccination are very important to determine the status of the vaccinated persons.”

If these figures are correct, this would be a global mass murder without equals. Considering that around five billion people worldwide were given such COVID vaccines, this would imply a number of 150 million deaths. And that would only be those deaths that took place within a very short time. What about those people who suffered myocarditis because of this “vaccine” and die as a result years later? What about those people whose immune system has been broken (VAIDS) and who die over time from other infectious diseases? What about all the cases of turbo cancer associated with the experimental gene injections?

It once again shows very clearly that more extensive studies need to be carried out on a global level in order to actually quantify the dangerousness of experimental COVID vaccines and injections of genetic sources. This study is another important piece of puzzle for creating an overall picture.